

Aivan lopuksi: anestesiaosastoilla tulisi olla so-
vittu toimintamalli tahattoman hereillä olon sat-
tuessa. Olisiko tässä SOP:n laatimisen paikka? ■

Viitteet

1. White DC. Anaesthesia: a privation of the senses. Kirjassa: Consciousness, Awareness and Pain in General Anaesthesia. 1. painos, s. 1-9. Toim. Rosen M, Lunn NJ. Butterworths, 1987.
2. Anonymous. Curare in Anaesthesia (editorial). The Lancet 1945; 246: 81-2.
3. Hutchinson R. Awareness during surgery. Br J Anaesth 1961; 33: 463-9.
4. Ranta S, Laurila R, Saario J, ym. Awareness with recall during general anesthesia: incidence and risk factors. Anesth Analg 1998; 86: 1084-9.
5. Sandin R, Enlund G, Samuelsson P, Lennmarken C. Awareness during anaesthesia: a prospective case study. The Lancet 2000; 355: 707-11.
6. Sebel P, Bowdle T, Ghoneim M, ym. The incidence of awareness during anesthesia: a multicenter United States study. Anesth Analg 2004; 99: 833-9.
7. Myles P, Leslie K, McNeil J, ym. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. Lancet 2004; 363: 1757-63.
8. Avidan M, Zhang L, Burnside B, ym. Anesthesia awareness and the bispectral index. N Engl J Med 2008; 358: 1097-108.
9. Avidan M, Jacobsohn E, Glick D, ym. Prevention of intraoperative awareness in a high-risk surgical population. N Engl J Med 2011; 365: 591-600.
10. Mashour G, Shanks A, Tremper K, ym. Prevention of intraoperative awareness with explicit recall in an unselected surgical population: A randomized comparative effectiveness trial. Anesthesiology 2012; 117: 717-25.
11. Aho AJ, Lyytikäinen LP, Yli-Hankala A, ym. Explaining Entropy responses after a noxious stimulus, with or without neuromuscular blocking agents, by means of the raw electroencephalographic and electromyographic characteristics. Br J Anaesth 2011; 106: 69-76.
12. Hart SM, Buchannan CR, Sleigh JW. A failure of M-Entropy to correctly detect burst suppression leading to sevoflurane overdosage. Anaesth Intensive Care 2009; 37: 1002-4.
13. Meyer B, Blacher R. A traumatic neurotic reaction induced by succinylcholine chloride. N Y State J Med 1961; 61: 1255-61.
14. Leslie K, Chan M, Myles P, ym. Post-traumatic stress disorder in aware patients from the B-aware trial. Anesth Analg 2010; 110: 823-8.

8th course of

1st announcement

Emergency Management of Severe Burns (EMSB)

15th April, 2016

Helsinki, Finland

Registration: sites.google.com/site/EMSBNordic

Information: timo.kukkonen@gmail.com

maarit.hult@hus.fi



Helsinki Burn Centre

Plastic Surgery and Burn Care Education
and Development Association